

Travelling waves and QCD saturation

Guillaume Beuf

CEA / DSM / SPhT

[From BK to FKPP](#)

[Solutions of the FKPP equation](#)

[Back to BK](#)

[BK with running coupling](#)

[BK at NLL](#)

An introduction to the traveling wave method in gluon saturation

S. Munier, R. Peschanski (2003-04)

- Link between saturation and nonlinear physics
- Nonlinear wave fronts formation and universal properties
- Results for the Balitsky-Kovchegov equation with fixed or running coupling

A more recent result

S. Sapeta, R. Peschanski (2006), G. B., R. Peschanski (2007)

- Stability with respect to higher orders

Balitsky-Kovchegov equation

From BK to FKPP

● Balitsky-Kovchegov equation

● 1D-BK equation

● Mapping to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

Balitsky (1996), Kovchegov (1999, 2000)

$$\begin{aligned} \partial_Y T_Y(\mathbf{x}, \mathbf{y}) = & \bar{\alpha} \int \frac{d^2 \mathbf{z}}{2\pi} \frac{(\mathbf{x} - \mathbf{y})^2}{(\mathbf{x} - \mathbf{z})^2 (\mathbf{z} - \mathbf{y})^2} [T_Y(\mathbf{x}, \mathbf{z}) \\ & + T_Y(\mathbf{z}, \mathbf{y}) - T_Y(\mathbf{x}, \mathbf{y}) - T_Y(\mathbf{x}, \mathbf{z}) T_Y(\mathbf{z}, \mathbf{y})] \end{aligned}$$

- BFKL kernel \Rightarrow exponential growth of $T_Y(\mathbf{x}, \mathbf{y})$
- Nonlinear damping \Rightarrow saturation at $T_Y(\mathbf{x}, \mathbf{y}) = 1$

1D-BK equation

From BK to FKPP

● Balitsky-Kovchegov equation

● 1D-BK equation

● Mapping to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- Fourier transform $\mathbf{x} - \mathbf{y} \mapsto \mathbf{k}$ and $(\mathbf{x} + \mathbf{y})/2 \mapsto \mathbf{q}$
- Restriction to zero momentum transfert $q = 0$
- Rotationnal invariance $\Rightarrow \tilde{T}_Y(\mathbf{k}, \mathbf{q} = 0) \equiv N(\log(k^2/Q_0^2), Y)$

$$\partial_Y N(L, Y) = \bar{\alpha} [\chi_{LL}(-\partial_L)N(L, Y) - N^2(L, Y)]$$

With $\chi_{LL}(\gamma) = 2\Psi(1) - \Psi(\gamma) - \Psi(1 - \gamma)$.

Munier, Peschanski (2003)

■ Diffusive approximation:

$$\chi_{LL}(-\partial_L) \simeq \chi_{LL}(\frac{1}{2}) + \frac{1}{2}\chi_{LL}''(\frac{1}{2}) \left(\frac{1}{2} + \partial_L\right)^2$$

■ Change of variables: $t \propto \bar{\alpha}Y$, $x = C_1L + C_2\bar{\alpha}Y$, and

$$u(t, x) \propto N(L, Y)$$

\Rightarrow FKPP equation


$$\partial_t u(t, x) = \partial_x^2 u(t, x) + u(t, x) - u^2(t, x)$$

Fisher (1937), Kolmogorov, Petrovsky, Piscounov (1937)

Uniformly translating fronts

$$u(t, x) = \phi_v(z) , \quad \text{with} \quad z \equiv x - vt$$

$$\begin{cases} -v \phi_v'(z) = \phi_v''(z) + \phi_v(z) - \phi_v^2(z) \\ \phi_v(-\infty) = 1 \quad \text{and} \quad \phi_v(+\infty) = 0 \end{cases}$$



From BK to FKPP

Solutions of the FKPP equation

● Uniformly translating fronts

● Uniformly translating fronts

● Uniformly translating fronts

● Generic initial condition

● Generic initial condition

● Asymptotic front

● Convergence to the asymptotic front

● Solution of the FKPP equation

● Universality

Back to BK

BK with running coupling

BK at NLL

Uniformly translating fronts

From BK to FKPP

Solutions of the FKPP equation

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- Universality

Back to BK


BK with running coupling

BK at NLL

Linear regime at large z :

$$\phi_v(z) \propto e^{-\gamma z} \quad \text{for } z \rightarrow \infty \quad \text{where } v = \gamma + \frac{1}{\gamma}$$

First case: $v \geq 2$




Uniformly translating fronts

Linear regime at large z :

$$\phi_v(z) \propto e^{-\gamma z} \quad \text{for } z \rightarrow \infty \quad \text{where } v = \gamma + \frac{1}{\gamma}$$

Second case: $v < 2$



From BK to FKPP

Solutions of the FKPP equation

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- Universality

Back to BK

BK with running coupling

BK at NLL

Uniformly translating fronts

From BK to FKPP

Solutions of the FKPP equation

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- Universality

Back to BK

BK with running coupling

BK at NLL

Linear regime at large z :

$$\phi_v(z) \propto e^{-\gamma z} \quad \text{for } z \rightarrow \infty \quad \text{where } v = \gamma + \frac{1}{\gamma}$$

Assumption: $u(t, x) \geq 0$

$$\Rightarrow v \geq 2$$

$$v = v(\gamma), \quad \text{for all } \gamma > 0$$

$$v_{min} = v(1) = 2$$

[From BK to FKPP](#)

[Solutions of the FKPP equation](#)

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- **Generic initial condition**
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- Universality


[Back to BK](#)

[BK with running coupling](#)

[BK at NLL](#)

Local study in the linear regime at large z :

If $v_1 < v_2$



[From BK to FKPP](#)

[Solutions of the FKPP equation](#)

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- **Generic initial condition**
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- Universality


[Back to BK](#)

[BK with running coupling](#)


[BK at NLL](#)

Local study in the linear regime at large z :

If $v_1 > v_2$



Asymptotic front



Nonlinear damping \Rightarrow flat solution at small x .

$$\text{For } t \rightarrow \infty, \quad u(t, x) \sim \begin{cases} e^{-\gamma_0 z} & \text{if } \gamma_0 < \gamma_c = 1 \\ e^{-\gamma_c z} & \text{if } \gamma_0 \geq \gamma_c \end{cases} \quad (1)$$

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- **Convergence to the asymptotic front**
- Solution of the FKPP equation
- Universality

Ansatz:

$$u(t, x) = t^\alpha G\left(\frac{\xi}{t^\alpha}\right) e^{-\gamma_c \xi}$$

$$\xi = x - v_c t + c(t)$$

for $t \rightarrow \infty$, and $\xi \leq \mathcal{O}(t^\alpha)$.

The FKPP equation gives then

$$\alpha = \frac{1}{2}, \quad \dot{c}(t) = \frac{\beta}{t}$$

$$0 = G''(z) + \frac{z}{2} G'(z) + \left(\beta - \frac{1}{2}\right) G(z)$$

Boundary conditions:

- $G(z)$ bounded for $z \rightarrow \infty$.
- $G(z) \sim z$ for $z \rightarrow 0$ (nonlinearities).

Solution of the FKPP equation

From BK to FKPP

Solutions of the FKPP equation

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- **Solution of the FKPP equation**
- Universality

Back to BK

BK with running coupling

BK at NLL

For $t \rightarrow \infty$, and $\mathcal{O}(1) \leq \xi \leq \mathcal{O}(\sqrt{t})$:

$$u(t, x) = A \xi e^{-\frac{\xi^2}{4t}} e^{-\xi}$$

$$\xi = x - 2t + \frac{3}{2} \log t$$

For $\xi \gg \sqrt{t}$: initial condition still relevant.

For $\xi \leq \mathcal{O}(1)$: nonlinear term relevant.

From BK to FKPP

Solutions of the FKPP equation

- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Uniformly translating fronts
- Generic initial condition
- Generic initial condition
- Asymptotic front
- Convergence to the asymptotic front
- Solution of the FKPP equation
- **Universality**

Back to BK

BK with running coupling

BK at NLL

For an FKPP-like equation with

- an unstable homogeneous equilibrium state
 - a family of uniformly translating front solutions
 - an effective nonlinear damping
 - a steep enough initial condition
- ⇒ Universal traveling wave asymptotic solution, independant of the precise form of
- the nonlinearities
 - the initial condition

Bramson (1983), Brunet, Derrida (1997), Ebert, van Saarloos (2000)

Solution in the linear regime:

$$N(L, Y) = \int \frac{d\gamma}{2\pi i} e^{-\gamma L + \bar{\alpha} \chi_{LL}(\gamma) Y} N_0(\gamma)$$

Dispersion relation:

$$v = \bar{\alpha} \frac{\chi_{LL}(\gamma)}{\gamma}$$

Critical parameters:

$$\chi_{LL}(\gamma_c) = \gamma_c \chi'_{LL}(\gamma_c) \Rightarrow \gamma_c = 0.6275\dots$$

$$v_c = \bar{\alpha} \frac{\chi_{LL}(\gamma_c)}{\gamma_c}$$

Initial condition:

$$\text{Color transparency} \Rightarrow N(L, Y) \propto e^{-L} \quad \text{for} \quad L \rightarrow \infty$$

Asymptotic solution of BK

Mueller, Triantafyllopoulos (2002), Munier, Peschanski (2003)

Universal asymptotic solution:

$$N(L, Y) = A \xi e^{-\frac{\xi^2}{2\bar{\alpha}\chi_{LL}''(\gamma_c)Y}} e^{-\gamma_c \xi}$$

$$\xi = L - \bar{\alpha} \frac{\chi_{LL}(\gamma_c)}{\gamma_c} Y + \frac{3}{2\gamma_c} \log Y ,$$

for $Y \rightarrow \infty$, and $\xi \leq \mathcal{O}(\sqrt{Y})$.

\Rightarrow Geometric scaling in $\tau = e^\xi = \frac{k^2}{Q_0^2} e^{-\bar{\alpha} \frac{\chi_{LL}(\gamma_c)}{\gamma_c} Y} Y^{\frac{3}{2\gamma_c}}$,

and scaling violations if $\xi \geq \mathcal{O}(\sqrt{Y})$.

A BK equation with running coupling

[From BK to FKPP](#)

[Solutions of the FKPP equation](#)

[Back to BK](#)

[BK with running coupling](#)

● [Equation with running coupling](#)

● [Approximate linear solution](#)

● [Dispersion relation](#)

● [Critical parameters](#)

● [Resolution](#)

[BK at NLL](#)

Choice: let us take $\bar{\alpha}$ at the scale k_T of the parent dipole.

$$\bar{\alpha}(k_T^2) = \frac{1}{b \log(k_T^2/\Lambda^2)} = \frac{1}{b L}$$

$$bL \partial_Y N(L, Y) = \chi_{LL}(-\partial_L) N(L, Y) - N^2(L, Y)$$

Approximate linear solution

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

- Equation with running coupling
- **Approximate linear solution**
- Dispersion relation
- Critical parameters
- Resolution

BK at NLL

$$N(L, Y) = \int \frac{d\gamma}{2\pi i} \int \frac{d\omega}{2\pi i} e^{-\gamma L + \omega Y + \frac{1}{b\omega}} X(\gamma) N_0(\gamma, \omega)$$

with

$$X(\gamma) = \int_{\hat{\gamma}}^{\gamma} d\gamma' \chi_{LL}(\gamma')$$

is an approximate solution of the linear equation at large L , because the large L saddle point equation is

$$L = \frac{1}{b\omega} \chi_{LL}(\gamma)$$

Dispersion relation

$$N(L, Y) = \int \frac{d\gamma}{2\pi i} \int \frac{d\omega}{2\pi i} e^{-\gamma L + \omega Y + \frac{1}{b\omega} X(\gamma)} N_0(\gamma, \omega)$$

At large Y , the saddle point approximation in ω gives

$$\omega_s = \sqrt{\frac{X(\gamma)}{bY}}$$

$$N(L, Y) \sim \int \frac{d\gamma}{2\pi i} e^{-\gamma L + \sqrt{\frac{4X(\gamma)}{b}} \sqrt{Y}} N_0(\gamma)$$

\Rightarrow dispersion relation at large Y and L :

$$v(\gamma) = \frac{1}{\gamma} \sqrt{\frac{4X(\gamma)}{b}}$$

The effective time for the wave is \sqrt{Y} and not Y .

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

- Equation with running coupling
- Approximate linear solution
- Dispersion relation
- Critical parameters
- Resolution

BK at NLL

$$v(\gamma) = \frac{1}{\gamma} \sqrt{\frac{4X(\gamma)}{b}}$$

$$X(\gamma) = \int_{\hat{\gamma}}^{\gamma} d\gamma' \chi_{LL}(\gamma')$$

$v_c = \min v(\gamma) = v(\gamma_c)$ depend on $\hat{\gamma}$. Let us choose $\hat{\gamma}$ such that

$$\frac{dv_c(\hat{\gamma})}{d\hat{\gamma}} = 0$$

⇒ Critical parameters:

$$\chi_{LL}(\gamma_c) = \gamma_c \chi'_{LL}(\gamma_c) \quad \gamma_c \simeq 0.6275$$

$$v_c = \sqrt{\frac{2\chi_{LL}(\gamma_c)}{b\gamma_c}}$$

- Equation with running coupling
- Approximate linear solution
- Dispersion relation
- Critical parameters
- Resolution

Expansion of the kernel around $\gamma \sim \gamma_c$

$$\frac{bL}{2\sqrt{Y}} \partial_{\sqrt{Y}} N = \left[-\frac{bv_c^2}{2} \partial_L + \frac{1}{2} \chi_{LL}''(\gamma_c) (\partial_L^2 + 2\gamma_c \partial_L + \gamma_c^2) + \dots \right] N$$

Then, using the same Ansatz:

$$\begin{aligned}
 N(L, Y) &= A Y^{1/6} \text{Ai} \left(\bar{\xi}_1 + \left(\frac{\sqrt{2b\gamma_c \chi_{LL}(\gamma_c)}}{\chi_{LL}''(\gamma_c)} \right)^{1/3} \frac{\xi}{Y^{1/6}} \right) e^{-\gamma_c \xi} \\
 \xi &\equiv \log \left(\frac{k^2}{Q_s^2(Y)} \right) \\
 &= L - \sqrt{\frac{2\chi_{LL}(\gamma_c)Y}{b\gamma_c}} - \frac{3\bar{\xi}_1}{4} \left(\frac{\chi_{LL}''(\gamma_c)}{\sqrt{2b\gamma_c \chi_{LL}(\gamma_c)}} \right)^{\frac{1}{3}} Y^{\frac{1}{6}} + \mathcal{O}(Y^{-\frac{1}{6}})
 \end{aligned}$$

General form:

$$\begin{aligned} \partial_Y N(L, Y) = & \bar{\alpha} [\chi_{LL}(-\partial_L) + \bar{\alpha} \chi_{NLL}(-\partial_L)] N(L, Y) \\ & - \bar{\alpha} [N^2(L, Y) + \bar{\alpha} (\text{NLL nonlinear terms})] \\ & + \bar{\alpha}^2 (\text{New terms ?}) \end{aligned}$$

Collecting the running coupling terms:

$$\begin{aligned} \partial_Y N(L, Y) = & \bar{\alpha}(L) \left[\chi_{LL}(-\partial_L) + \bar{\alpha}(L) \chi^{(1)}(-\partial_L) \right] N(L, Y) \\ & - \bar{\alpha}(L) (\text{Nonlinear terms}) \\ & + \bar{\alpha}^2(L) (\text{New terms ?}) \end{aligned}$$

Effective linear equation

Linear part of the NLL equation:

$$\partial_Y N(L, Y) = \frac{1}{bL} \left[\chi_{LL}(-\partial_L) + \frac{1}{bL} \chi^{(1)}(-\partial_L) \right] N(L, Y)$$

Is it equivalent to an effective equation

$$\partial_Y N(L, Y) = \frac{1}{bL} \kappa(-\partial_L, \partial_Y) N(L, Y) ?$$

Ciafaloni, Colferai (1998)

- From BK to FKPP
- Solutions of the FKPP equation
- Back to BK
- BK with running coupling
- BK at NLL
 - BK at NLL
 - **Effective linear equation**
 - Effective linear equation
 - Omega expansion
 - Omega expansion
 - Approximate linear solution
 - Large Y saddle point
 - Critical parameters
 - Expansion of the effective equation
 - Result at NLL

[From BK to FKPP](#)

[Solutions of the FKPP equation](#)

[Back to BK](#)

[BK with running coupling](#)

[BK at NLL](#)

- BK at NLL
- Effective linear equation
- **Effective linear equation**
- Omega expansion
- Omega expansion
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL

In Laplace (or Mellin) space:

$$-\partial_L \rightarrow \gamma \quad \text{and} \quad \partial_Y \rightarrow \omega$$

$$\omega = \frac{1}{bL} \left[\chi_{LL}(\gamma) + \frac{1}{bL} \chi^{(1)}(\gamma) \right]$$

$$\omega = \frac{1}{bL} \kappa(\gamma, \omega)$$

They are equivalent if

$$\kappa(\gamma, \omega) = \chi_{LL}(\gamma) + \frac{\omega \chi^{(1)}(\gamma)}{\kappa(\gamma, \omega)}.$$

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- BK at NLL
- Effective linear equation
- Effective linear equation
- **Omega expansion**
- Omega expansion
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL

Effective kernel:

$$\begin{aligned}\kappa(\gamma, \omega) &= \chi_{LL}(\gamma) \frac{1}{2} \left[1 + \sqrt{1 + \frac{4\omega \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)}} \right] \\ &\simeq \chi_{LL}(\gamma) \left[1 + \frac{\omega \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} - \left(\frac{\omega \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} \right)^2 + \dots \right]\end{aligned}$$

The effective equation can be trusted if

$$\frac{\omega \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} \ll 1$$

Omega expansion

The LL and NLL kernel eigenfunctions have singularities in $\gamma \rightarrow 0$ (or $1 - \gamma \rightarrow 0$)

$$\chi_{LL}(\gamma) \propto \gamma^{-1} \quad (\text{or } (1 - \gamma)^{-1})$$

$$\chi^{(1)}(\gamma) \propto \gamma^{-3} \quad (\text{or } (1 - \gamma)^{-3})$$

$$\frac{\omega \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} \propto \frac{\omega}{\gamma} \quad \left(\text{or } \frac{\omega}{1 - \gamma} \right)$$

\Rightarrow The effective equation is valid if $\omega \ll \gamma, 1 - \gamma$

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- BK at NLL
- Effective linear equation
- Effective linear equation
- Omega expansion
- **Omega expansion**
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL

Approximate linear solution

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- BK at NLL
- Effective linear equation
- Effective linear equation
- Omega expansion
- Omega expansion
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL

$$N(L, Y) = \int \frac{d\gamma}{2\pi i} \int \frac{d\omega}{2\pi i} e^{-\gamma L + \omega Y + \frac{1}{b\omega}} X(\gamma, \omega) N_0(\gamma, \omega)$$

with

$$X(\gamma, \omega) = \int_{\hat{\gamma}}^{\gamma} d\gamma' \kappa(\gamma', \omega)$$

is an approximate solution of the linear equation at large L , because the large L saddle point equation is

$$L = \frac{1}{b\omega} \kappa(\gamma, \omega)$$

which is equivalent to

$$\omega = \frac{1}{bL} \left[\chi_{LL}(\gamma) + \frac{1}{bL} \chi^{(1)}(\gamma) \right] \quad \text{if } \omega \ll \gamma, 1 - \gamma.$$

Large Y saddle point

$$N(L, Y) = \int \frac{d\gamma}{2\pi i} \int \frac{d\omega}{2\pi i} e^{-\gamma L + \omega Y + \frac{1}{b\omega}} X(\gamma, \omega) N_0(\gamma, \omega)$$

At large Y , the saddle point approximation in ω gives

$$\begin{aligned} Y b \omega_s^2 &= X(\gamma, \omega_s) - \omega_s \dot{X}(\gamma, \omega_s) \\ &= \int_{\hat{\gamma}}^{\gamma} d\gamma' \chi_{LL}(\gamma') \left[1 + \left(\frac{\omega_s \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} \right)^2 \right. \\ &\quad \left. - 4 \left(\frac{\omega_s \chi^{(1)}(\gamma)}{\chi_{LL}^2(\gamma)} \right)^3 + \dots \right] \end{aligned}$$

From BK to FKPP
Solutions of the FKPP equation
Back to BK
BK with running coupling
BK at NLL
● BK at NLL
● Effective linear equation
● Effective linear equation
● Omega expansion
● Omega expansion
● Approximate linear solution
● Large Y saddle point
● Critical parameters
● Expansion of the effective equation
● Result at NLL

Critical parameters

If $\gamma \rightarrow 0$:

$$Y b \omega_s^2 = \log \gamma + \text{constant} - \frac{1}{2} \left(\frac{C \omega_s}{\gamma} \right)^2 + \frac{4}{3} \left(\frac{C \omega_s}{\gamma} \right)^3 + \dots$$

For large Y , $\omega_s \propto Y^{-1/2}$, and the higher orders are suppressed if $\omega_s \ll \gamma, 1 - \gamma$.

\Rightarrow same γ_c and v_c as in the simplest equation with running coupling. And the convergence can start when

$$Y > \gamma_c^{-2}, (1 - \gamma_c)^{-2}$$

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- BK at NLL
- Effective linear equation
- Effective linear equation
- Omega expansion
- Omega expansion
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL

Expanding the effective kernel around $\gamma \sim \gamma_c$ and $\omega \sim 0$:

$$\begin{aligned} \frac{bL}{2\sqrt{Y}} \partial_{\sqrt{Y}} N = & \left(-\frac{bv_c^2}{2} \partial_L + \frac{1}{2} \chi_{LL}''(\gamma_c)(\partial_L^2 + 2\gamma_c \partial_L + \gamma_c^2) + \dots \right. \\ & + \frac{1}{2\sqrt{Y}} \frac{\chi^{(1)}(\gamma_c)}{\chi_{LL}(\gamma_c)} \partial_{\sqrt{Y}} - \frac{1}{2\sqrt{Y}} \left(\partial_\gamma \frac{\chi^{(1)}(\gamma)}{\chi_{LL}(\gamma)} \right)_{\gamma=\gamma_c} (\gamma_c + \partial_L) \partial_{\sqrt{Y}} \\ & \left. + \dots \right) N \end{aligned}$$

The new terms, of order $Y^{-1/2}$ doesn't contribute to the universal subasymptotic behavior.

At large enough rapidity ($Y^{-1/2} \ll 1$), the solution of the full NLL BK equation in the geometric scaling region converge to the solution of the LL BK equation with running coupling. Then, they converge to their asymptotic solution.

$$\begin{aligned}
 N(L, Y) &= A Y^{1/6} \text{Ai} \left(\bar{\xi}_1 + \left(\frac{\sqrt{2b\gamma_c \chi_{LL}(\gamma_c)}}{\chi_{LL}''(\gamma_c)} \right)^{1/3} \frac{\xi}{Y^{1/6}} \right) e^{-\gamma_c \xi} \\
 \xi &\equiv \log \left(\frac{k^2}{Q_s^2(Y)} \right) \\
 &= L - \sqrt{\frac{2\chi_{LL}(\gamma_c)Y}{b\gamma_c}} - \frac{3\bar{\xi}_1}{4} \left(\frac{\chi_{LL}''(\gamma_c)}{\sqrt{2b\gamma_c \chi_{LL}(\gamma_c)}} \right)^{\frac{1}{3}} Y^{\frac{1}{6}}
 \end{aligned}$$

From BK to FKPP

Solutions of the FKPP equation

Back to BK

BK with running coupling

BK at NLL

- BK at NLL
- Effective linear equation
- Effective linear equation
- Omega expansion
- Omega expansion
- Approximate linear solution
- Large Y saddle point
- Critical parameters
- Expansion of the effective equation
- Result at NLL